arXiv:1607.04111v1 [math.DG] 14 Jul 2016

GENERAL ROTATIONAL SURFACES IN PSEUDO-EUCLIDEAN 4-SPACE WITH NEUTRAL METRIC

YANA ALEKSIEVA, VELICHKA MILOUSHEVA, NURETTIN CENK TURGAY

Abstract. We define general rotational surfaces of elliptic and hyperbolic type in the pseudo-Euclidean 4-space with neutral metric which are analogous to the general rotational surfaces of C. Moore in the Euclidean 4-space. We study Lorentz general rotational surfaces with plane meridian curves and give the complete classification of minimal general rotational surfaces of elliptic and hyperbolic type, general rotational surfaces with parallel normalized mean curvature vector field, flat general rotational surfaces, and general rotational surfaces with flat normal connection.

1. Introduction

In [14] C. Moore introduced general rotational surfaces in the 4-dimensional Euclidean space $\mathbb{E}^4$ and described a special case of general rotational surfaces with constant Gauss curvature [15]. In [13] the second author studied general rotational surfaces whose meridians lie in two-dimensional planes and completely classified all minimal super-conformal general rotational surfaces in $\mathbb{E}^4$. The minimal non-super-conformal general rotational surfaces with plane meridian curves are described by U. Dursun and the third author in [8]. In [9] the complete classification of general rotational surfaces consisting of parabolic points is given. General rotational surfaces with plane meridian curves and pointwise 1-type Gauss map are studied in [7].

In [11] G. Ganchev and the second author studied spacelike general rotational surfaces in the 4-dimensional Minkowski space $\mathbb{E}_1^4$ that are analogous to the general rotational surfaces of C. Moore in $\mathbb{E}^4$ and described analytically flat general rotation surfaces and general rotational surfaces with flat normal connection. The classification of minimal general rotational surfaces in $\mathbb{E}_1^4$ and general rotational surfaces consisting of parabolic points is also given in [11]. Spacelike general rotational surfaces in $\mathbb{E}_1^4$ with meridian curves lying in 2-dimensional planes and having pointwise 1-type Gauss map are studied in [6].

In the present paper we define general rotational surfaces of elliptic and hyperbolic type in the pseudo-Euclidean 4-space with neutral metric $\mathbb{E}_2^4$ which are analogous to the general rotational surfaces in $\mathbb{E}^4$ and $\mathbb{E}_1^4$. We study Lorentz general rotational surfaces with meridian curves lying in 2-dimensional planes. In Theorem 4.1 and Theorem 4.2 we give the complete classification of minimal general rotational surfaces of elliptic and hyperbolic type. Theorem 5.1 and Theorem 5.3 classify general rotational surfaces of elliptic and hyperbolic type with parallel normalized mean curvature vector field. The classification of flat general rotational surfaces of elliptic and hyperbolic type is given in Theorem 6.1 and Theorem 6.2, respectively. In the last section we describe all general rotational surfaces of elliptic and hyperbolic type with flat normal connection (Theorem 7.1 and Theorem 7.2).

2010 *Mathematics Subject Classification.* Primary 53B30, Secondary 53A35, 53B25.

Key words and phrases. Pseudo-Euclidean space, Lorentz surfaces, general rotational surfaces, minimal surfaces, parallel mean curvature vector.

2. Preliminaries

Let $\mathbb{E}_2^4$ be the pseudo-Euclidean 4-space endowed with the canonical pseudo-Euclidean metric of index 2 given by

$$g_0 = dx_1^2 + dx_2^2 - dx_3^2 - dx_4^2,$$

where (x_1, x_2, x_3, x_4) is a rectangular coordinate system of $\mathbb{E}_2^4$. As usual, we denote by $\langle .\,, . \rangle$ the indefinite inner scalar product with respect to g_0. A non-zero vector v is called *spacelike* (respectively, *timelike*) if $\langle v, v\rangle > 0$ (respectively, $\langle v, v\rangle < 0$). A vector v is called *lightlike* if it is nonzero and satisfies $\langle v, v\rangle = 0$.

A surface M_1^2 in $\mathbb{E}_2^4$ is called *Lorentz* if the induced metric g on M_1^2 is Lorentzian, i.e. at each point $p \in M_1^2$ we have the following decomposition

$$\mathbb{E}_2^4 = T_pM_1^2 \oplus N_pM_1^2$$

with the property that the restriction of the metric onto the tangent space $T_pM_1^2$ is of signature $(1, 1)$, and the restriction of the metric onto the normal space $N_pM_1^2$ is of signature $(1, 1)$.

Denote by ∇ and ∇' the Levi Civita connections of M_1^2 and $\mathbb{E}_2^4$, respectively. Let x and y be vector fields tangent to M_1^2 and ξ be a normal vector field. The formulas of Gauss and Weingarten are given respectively by

$$\nabla'_x y = \nabla_x y + \sigma(x, y);$$
$$\nabla'_x \xi = -A_\xi x + D_x \xi,$$

where σ is the second fundamental form, D is the normal connection, and A_ξ is the shape operator with respect to ξ. In general, A_ξ is not diagonalizable.

The mean curvature vector field H of M_1^2 is defined as $H = \dfrac{1}{2}\operatorname{tr}\sigma$. A surface M_1^2 is called *minimal* if its mean curvature vector vanishes identically, i.e. $H = 0$. A surface M_1^2 is called *quasi-minimal* (or *pseudo-minimal*) if its mean curvature vector is lightlike at each point, i.e. $H \neq 0$ and $\langle H, H\rangle = 0$ [16].

A normal vector field ξ on M_1^2 is called *parallel in the normal bundle* (or simply *parallel*) if $D\xi = 0$ holds identically [5]. The surface M_1^2 is said to have *parallel mean curvature vector field* if its mean curvature vector H is parallel, i.e. $DH = 0$.

A natural extension of the class of surfaces with parallel mean curvature vector field are surfaces with parallel normalized mean curvature vector field. A surface M_1^2 is said to have *parallel normalized mean curvature vector field* if the mean curvature vector H is non-zero and there exists a unit vector field in the direction of the mean curvature vector field which is parallel in the normal bundle [4]. It is easy to see that if M_1^2 is a surface with non-zero parallel mean curvature vector field H (i.e. $DH = 0$), then M_1^2 is a surface with parallel normalized mean curvature vector field, but the converse is not true in general. It is true only in the case $\|H\| = const$.

3. General Rotational Surfaces of Elliptic and Hyperbolic Type

General rotational surfaces in the Euclidean 4-space $\mathbb{E}^4$ were introduced by C. Moore [14] as follows. Let $m : x(u) = \big(x^1(u), x^2(u), x^3(u), x^4(u)\big)$; $u \in J \subset \mathbb{R}$ be a smooth curve in $\mathbb{E}^4$, and α, β be constants. A general rotation of the meridian curve m is defined by

$$X(u, v) = \big(X^1(u, v), X^2(u, v), X^3(u, v), X^4(u, v)\big)\,,$$

where

$$X^1(u,v) = x^1(u)\cos\alpha v - x^2(u)\sin\alpha v; \qquad X^3(u,v) = x^3(u)\cos\beta v - x^4(u)\sin\beta v;$$
$$X^2(u,v) = x^1(u)\sin\alpha v + x^2(u)\cos\alpha v; \qquad X^4(u,v) = x^3(u)\sin\beta v + x^4(u)\cos\beta v.$$

If $\beta = 0$, $x^2(u) = 0$ the plane Oe_3e_4 is fixed and one gets the classical rotation about a fixed two-dimensional axis.

Similarly to the general rotations in the Euclidean space $\mathbb{E}^4$ one can consider general rotational surfaces in the Minkowski 4-space $\mathbb{E}_1^4$ (see [11] and [6]).

Now we shall define general rotational surfaces of Moore type in the pseudo-Euclidean 4-space $\mathbb{E}_2^4$.

Let $Oe_1e_2e_3e_4$ be an orthonormal base of $\mathbb{E}_2^4$, such that $\langle e_1, e_1\rangle = \langle e_2, e_2\rangle = 1$, and $\langle e_3, e_3\rangle = \langle e_4, e_4\rangle = -1$. Let $m : x(u) = \big(x^1(u), x^2(u), x^3(u), x^4(u)\big)$; $u \in J \subset \mathbb{R}$ be a smooth spacelike or timelike curve in $\mathbb{E}_2^4$, and α, β be constants. A general rotational surface of elliptic type can be defined as follows:

$$X(u,v) = \big(X^1(u,v), X^2(u,v), X^3(u,v), X^4(u,v)\big),$$

where

$$\begin{aligned} &X^1(u,v) = x^1(u)\cos\alpha v - x^2(u)\sin\alpha v; \qquad X^3(u,v) = x^3(u)\cos\beta v - x^4(u)\sin\beta v;\\ &X^2(u,v) = x^1(u)\sin\alpha v + x^2(u)\cos\alpha v; \qquad X^4(u,v) = x^3(u)\sin\beta v + x^4(u)\cos\beta v. \end{aligned} \tag{1}$$

If $\beta = 0$, $x^2(u) = 0$ one gets the surface with parametrization

$$X(u,v) = \big(x^1(u)\cos\alpha v, x^1(u)\sin\alpha v, x^3(u), x^4(u)\big),$$

which is a standard rotation of elliptic type about the 2-dimensional axis Oe_3e_4. In the case $\alpha = 0$, $x^4(u) = 0$ we get the surface

$$X(u,v) = \big(x^1(u), x^2(u), x^3(u)\cos\beta v, x^3(u)\sin\beta v\big),$$

which is a standard rotation of elliptic type about Oe_1e_2. If $\alpha > 0$ and $\beta > 0$ the surface defined by (1) is analogous to the general rotational surface of C. Moore in $\mathbb{E}^4$.

In the present paper we shall consider Lorentz general rotational surfaces of elliptic type for which $\alpha > 0$, $\beta > 0$, $x^2(u) = x^4(u) = 0$. In this case the meridian curve m lies in two-dimensional plane.

Similarly to the general rotational surfaces of elliptic type we define general rotational surfaces of hyperbolic type in $\mathbb{E}_2^4$ as follows:

$$\begin{aligned} &X^1(u,v) = x^1(u)\cosh\alpha v + x^3(u)\sinh\alpha v; \qquad X^3(u,v) = x^1(u)\sinh\alpha v + x^3(u)\cosh\alpha v;\\ &X^2(u,v) = x^2(u)\cosh\beta v + x^4(u)\sinh\beta v; \qquad X^4(u,v) = x^2(u)\sinh\beta v + x^4(u)\cosh\beta v. \end{aligned} \tag{2}$$

In the case $\beta = 0$, $x^3(u) = 0$ we obtain the surface with parametrization

$$X(u,v) = \big(x^1(u)\cosh\alpha v, x^2(u), x^1(u)\sinh\alpha v, x^4(u)\big),$$

which is a standard rotation of hyperbolic type about Oe_2e_4. In the case $\alpha = 0$, $x^4(u) = 0$ we get the surface

$$X(u,v) = \big(x^1(u), x^2(u)\cosh\beta v, x^3(u), x^2(u)\sinh\beta v\big),$$

which is a standard rotation of hyperbolic type about Oe_1e_3. If $\alpha > 0$ and $\beta > 0$ the surface defined by (2) is a general rotational surface of hyperbolic type in $\mathbb{E}_2^4$. We shall consider Lorentz general rotational surfaces of hyperbolic type for which $\alpha > 0$, $\beta > 0$, $x^3(u) = x^4(u) = 0$.

3.1. **General rotational surfaces of elliptic type with plane meridians.** Let $\mathcal{M}_1$ be the general rotational surface of elliptic type defined by:

$$\mathcal{M}_1 : z(u,v) = (f(u)\cos\alpha v, f(u)\sin\alpha v, g(u)\cos\beta v, g(u)\sin\beta v)\,, \tag{3}$$

where $u \in J \subset \mathbb{R}$, $v \in [0; 2\pi)$, $f(u)$ and $g(u)$ are smooth functions satisfying $\alpha^2 f^2(u) - \beta^2 g^2(u) < 0$, $f'^2(u) - g'^2(u) > 0$, and α, β are positive constants.

The tangent frame field $T_p\mathcal{M}_1$ is determined by the vector fields

$$\begin{aligned} z_u &= (f'(u)\cos\alpha v, f'(u)\sin\alpha v, g'(u)\cos\beta v, g'(u)\sin\beta v)\,;\\ z_v &= (-\alpha f(u)\sin\alpha v, \alpha f(u)\cos\alpha v, -\beta g(u)\sin\beta v, \beta g(u)\cos\beta v)\,. \end{aligned}$$

The coefficients of the first fundamental form of $\mathcal{M}_1$ are expressed by:

$$\begin{aligned} E &= \langle z_u, z_u\rangle = f'^2(u) - g'^2(v) > 0;\\ F &= \langle z_u, z_v\rangle = 0;\\ G &= \langle z_v, z_v\rangle = \alpha^2 f^2(u) - \beta^2 g^2(u) < 0. \end{aligned}$$

So, $\mathcal{M}_1$ is a Lorentz surface in $\mathbb{E}_2^4$.

We consider the following orthonormal tangent frame field:

$$x = \frac{z_u}{\sqrt{E}}; \quad y = \frac{z_v}{\sqrt{-G}},$$

which satisfy $\langle x, x\rangle = 1$, $\langle y, y\rangle = -1$, and $\langle x, y\rangle = 0$. Let n_1 and n_2 be the normal vector fields defined by:

$$\begin{aligned} n_1 &= \frac{1}{\sqrt{-\alpha^2 f^2(u) + \beta^2 g^2(u)}}\,(\beta g(u)\sin\alpha v, -\beta g(u)\cos\alpha v, \alpha f(u)\sin\beta v, -\alpha f(u)\cos\beta v)\,;\\ n_2 &= \frac{1}{\sqrt{f'^2(u) - g'^2(v)}}\,(g'(u)\cos\alpha v, g'(u)\sin\alpha v, f'(u)\cos\beta v, f'(u)\sin\beta v)\,. \end{aligned}$$

Note that $\langle n_1, n_1\rangle = 1$, $\langle n_2, n_2\rangle = -1$, $\langle n_1, n_2\rangle = 0$. Calculating the second derivatives

$$\begin{aligned} z_{uu} &= (f''(u)\cos\alpha v, f''(u)\sin\alpha v, g''(u)\cos\beta v, g''(u)\sin\beta v)\,;\\ z_{uv} &= (-\alpha f'(u)\sin\alpha v, \alpha f'(u)\cos\alpha v, -\beta g'(u)\sin\beta v, \beta g'(u)\cos\beta v)\,;\\ z_{vv} &= \big(-\alpha^2 f(u)\cos\alpha v, -\alpha^2 f(u)\sin\alpha v, -\beta^2 g(u)\cos\beta v, -\beta^2 g(u)\sin\beta v\big)\,, \end{aligned}$$

we obtain the following components of the second fundamental tensor:

$$\begin{aligned} &\langle z_{uu}, n_1\rangle = 0; && \langle z_{uu}, n_2\rangle = \frac{f''(u)g'(u) - g''(u)f'(u)}{\sqrt{f'^2(u) - g'^2(u)}};\\ &\langle z_{uv}, n_1\rangle = \frac{\alpha\beta(f(u)g'(u) - f'(u)g(u))}{\sqrt{\beta^2 g^2(u) - \alpha^2 f^2(u)}}; && \langle z_{uv}, n_2\rangle = 0;\\ &\langle z_{vv}, n_1\rangle = 0; && \langle z_{vv}, n_2\rangle = \frac{\beta^2 f'(u)g(u) - \alpha^2 f(u)g'(u))}{\sqrt{f'^2(u) - g'^2(u)}}. \end{aligned}$$

The above formulas imply

$$
\begin{aligned}
\sigma(x,x) &= \frac{f'(u)g''(u)-g'(u)f''(u)}{(\sqrt{f'^2(u)-g'^2(u)})^3}\, n_2;\\
\sigma(x,y) &= \frac{\alpha\beta(f(u)g'(u)-f'(u)g(u))}{\sqrt{f'^2(u)-g'^2(u)}(\beta^2g^2(u)-\alpha^2f^2(u))}\, n_1;\\
\sigma(y,y) &= \frac{(\alpha^2f(u)g'(u)-\beta^2f'(u)g(u))}{\sqrt{f'^2(u)-g'^2(u)}(\beta^2g^2(u)-\alpha^2f^2(u))}\, n_2.
\end{aligned}
\tag{4}
$$

Using that the Gauss curvature K is determined by the formula

$$K=\frac{\langle\sigma(x,x),\sigma(y,y)\rangle-\langle\sigma(x,y),\sigma(x,y)\rangle}{\langle x,x\rangle\langle y,y\rangle-\langle x,y\rangle^2},$$

we obtain

$$K=\frac{\alpha^2\beta^2(f'^2-g'^2)(fg'-f'g)^2-(\beta^2g^2-\alpha^2f^2)(\beta^2f'g-\alpha^2fg')(f'g''-f''g')}{(f'^2-g'^2)^2(\beta^2g^2-\alpha^2f^2)^2}.$$

Calculating the derivatives of the normal vector fields n_1 and n_2 we get

$$
\begin{aligned}
&D_xn_1=0; && D_xn_2=0;\\
&D_yn_1=\frac{\alpha\beta(ff'-gg')}{\sqrt{f'^2-g'^2}(\beta^2g^2-\alpha^2f^2)}\, n_2; && D_yn_2=\frac{\alpha\beta(ff'-gg')}{\sqrt{f'^2-g'^2}(\beta^2g^2-\alpha^2f^2)}\, n_1.
\end{aligned}
\tag{5}
$$

For the tangent vector fields x and y we obtain the following derivative formulas:

$$
\begin{aligned}
&\nabla_xx=0; && \nabla_xy=0;\\
&\nabla_yx=-\frac{\alpha^2ff'-\beta^2gg'}{\sqrt{f'^2-g'^2}(\beta^2g^2-\alpha^2f^2)}\, y; && \nabla_yy=-\frac{\alpha^2ff'-\beta^2gg'}{\sqrt{f'^2-g'^2}(\beta^2g^2-\alpha^2f^2)}\, x.
\end{aligned}
\tag{6}
$$

The curvature of the normal connection $\varkappa$ of $\mathcal{M}_1$ is determined by

$$\varkappa=\frac{\langle R^\perp(x,y)n_1,n_2\rangle}{\langle x,x\rangle\langle y,y\rangle-\langle x,y\rangle^2}=-\langle D_xD_yn_1-D_yD_xn_1-D_{[x,y]}n_1,n_2\rangle.$$

Formulas (5) and (6) imply that the curvature of the normal connection is given by the following expression:

$$\varkappa=\frac{-\alpha\beta(fg'-gf')\left((\beta^2g^2-\alpha^2f^2)(g'f''-f'g'')+(f'^2-g'^2)(\beta^2gf'-\alpha^2fg')\right)}{(f'^2-g'^2)^2(\beta^2g^2-\alpha^2f^2)^2}.\tag{7}$$

The normal mean curvature vector field H is defined by the formula

$$H=\frac{\sigma(x,x)-\sigma(y,y)}{2}.$$

Using the expressions for $\sigma(x,x)$ and $\sigma(y,y)$ in formulas (4) we get:

$$H=\frac{(f'^2-g'^2)(\beta^2gf'-\alpha^2fg')-(\beta^2g^2-\alpha^2f^2)(f''g'-f'g'')}{2\,(f'^2-g'^2)^{\frac{3}{2}}\,(\beta^2g^2-\alpha^2f^2)}\, n_2.\tag{8}$$

Formula (8) shows that in the case $H\neq0$ the normalized mean curvature vector field of a general rotational surface of elliptic type with plane meridian curves is timelike. So, we can formulate the following statement:

Proposition 3.1. *There are no quasi-minimal general rotational surfaces of elliptic type with plane meridian curves.*

Remark 3.2. Let us note that in the class of standard rotational surfaces in $\mathbb{E}^4_2$ there exist quasi-minimal surfaces and they are described in [10].

In our further considerations we shall use the following notations:

$$
\begin{aligned}
&\nu_1 = \frac{g'f'' - f'g''}{(f'^2 - g'^2)^{\frac{3}{2}}}; && \nu_2 = \frac{\beta^2 g f' - \alpha^2 f g'}{\sqrt{f'^2 - g'^2}(\beta^2 g^2 - \alpha^2 f^2)};\\
&\mu = \frac{\alpha\beta(fg' - gf')}{\sqrt{(f'^2 - g'^2)}(\beta^2 g^2 - \alpha^2 f^2)}; && \gamma_2 = \frac{\alpha^2 f f' - \beta^2 g g'}{\sqrt{f'^2 - g'^2}(\beta^2 g^2 - \alpha^2 f^2)};\\
&\beta_2 = \frac{\alpha\beta(ff' - gg')}{\sqrt{f'^2 - g'^2}(\beta^2 g^2 - \alpha^2 f^2)}.
\end{aligned}
\tag{9}
$$

Thus, equalities (4), (5), and (6) give the following derivative formulas of $\mathcal{M}_1$:

$$
\begin{aligned}
&\nabla'_x x = -\nu_1\, n_2; && \nabla'_x n_1 = \mu\, y;\\
&\nabla'_x y = \mu\, n_1; && \nabla'_y n_1 = -\mu\, x + \beta_2\, n_2;\\
&\nabla'_y x = -\gamma_2\, y + \mu\, n_1; && \nabla'_x n_2 = -\nu_1\, x;\\
&\nabla'_y y = -\gamma_2\, x - \nu_2\, n_2; && \nabla'_y n_2 = \nu_2\, y + \beta_2\, n_1.
\end{aligned}
\tag{10}
$$

Remark 3.3. In [2], an invariant local theory of Lorentz surfaces in the pseudo-Euclidean space $\mathbb{E}^4_2$ is developed and a family of eight geometric functions is introduced. It is proved that these geometric functions determine the surface up to a rigid motion in $\mathbb{E}^4_2$. The functions ν_1, ν_2, μ, γ_2, β_2 given in (9) are the geometric functions of the general rotational surface $\mathcal{M}_1$ in the sense of [2] (note that the other three geometric functions of $\mathcal{M}_1$ are equal to zero).

3.2. **General rotational surfaces of hyperbolic type with plane meridians.** Now we shall consider general rotational surfaces of hyperbolic type with plane meridian curves. Let $\mathcal{M}_2$ be the surface defined by

$$
\mathcal{M}_2 : z(u,v) = (f(u)\cosh\alpha v, g(u)\cosh\beta v, f(u)\sinh\alpha v, g(u)\sinh\beta v)\,,
\tag{11}
$$

where $u \in J \subset \mathbb{R}$, $v \in [0; 2\pi)$, $f(u)$ and $g(u)$ are smooth functions, which satisfy the conditions $\alpha^2 f^2(u) + \beta^2 g^2(u) > 0$, $f'^2(u) + g'^2(u) > 0$, $\alpha = const > 0$, $\beta = const > 0$.

The tangent vector fields z_u and z_v of $\mathcal{M}_2$ are:

$$
\begin{aligned}
z_u &= (f'(u)\cosh\alpha v, g'(u)\cosh\beta v, f'(u)\sinh\alpha v, g'(u)\sinh\beta v)\,,\\
z_v &= (\alpha f(u)\sinh\alpha v, \beta g(u)\sinh\beta v, \alpha f(u)\cosh\alpha v, \beta g(u)\cosh\beta v)\,.
\end{aligned}
$$

The coefficients of the first fundamental form of $\mathcal{M}_2$ are given by:

$$
E = f'^2(u) + g'^2(v), \quad F = 0, \quad G = -(\alpha^2 f^2 + \beta^2 g^2).
$$

So, $\mathcal{M}_2$ is a Lorentz surface.

Now we consider the following orthonormal tangent frame field $x = \dfrac{z_u}{\sqrt{E}}$, $y = \dfrac{z_v}{\sqrt{-G}}$, which satisfies $\langle x, x\rangle = 1$, $\langle y, y\rangle = -1$, $\langle x, y\rangle = 0$. We choose the following normal vector

fields of $\mathcal{M}_2$:

$$n_1 = \frac{1}{\sqrt{f'^2(u)+g'^2(v)}}\,(g'(u)\cosh\alpha v, -f'(u)\cosh\beta v, g'(u)\sinh\alpha v, -f'(u)\sinh\beta v)\,;$$
$$n_2 = \frac{1}{\sqrt{\alpha^2 f^2(u)+\beta^2 g^2(u)}}\,(\beta g(u)\sinh\alpha v, -\alpha f(u)\sinh\beta v, \beta g(u)\cosh\alpha v, -\alpha f(u)\cosh\beta v)\,,$$

for which we have $\langle n_1, n_1\rangle = 1$, $\langle n_2, n_2\rangle = -1$, $\langle n_1, n_2\rangle = 0$.

With respect to the frame field $\{x, y, n_1, n_2\}$ introduced above we obtain the following derivative formulas of $\mathcal{M}_2$:

$$\begin{aligned}
&\nabla'_x x = \nu_1\, n_1; && \nabla'_x n_1 = -\nu_1\, x;\\
&\nabla'_x y = -\mu\, n_2; && \nabla'_y n_1 = \nu_2\, y - \beta_2\, n_2;\\
&\nabla'_y x = -\gamma_2\, y - \mu\, n_2; && \nabla'_x n_2 = \mu\, y;\\
&\nabla'_y y = -\gamma_2\, x + \nu_2\, n_1; && \nabla'_y n_2 = -\mu\, x - \beta_2\, n_1,
\end{aligned} \tag{12}$$

where the functions ν_1, ν_2, μ, γ_2, β_2 are expressed by:

$$\begin{aligned}
&\nu_1 = \frac{f''g' - f'g''}{\left(\sqrt{f'^2+g'^2}\right)^3}; && \nu_2 = \frac{\alpha^2 fg' - \beta^2 gf'}{\sqrt{f'^2+g'^2}(\alpha^2 f^2+\beta^2 g^2)};\\
&\mu = \frac{\alpha\beta(fg'-f'g)}{\sqrt{f'^2+g'^2}(\alpha^2 f^2+\beta^2 g^2)}; && \gamma_2 = -\frac{\alpha^2 ff' + \beta^2 gg'}{\sqrt{f'^2+g'^2}(\alpha^2 f^2+\beta^2 g^2)};\\
&\beta_2 = -\frac{\alpha\beta(ff'+gg')}{\sqrt{f'^2+g'^2}(\alpha^2 f^2+\beta^2 g^2)}.
\end{aligned} \tag{13}$$

These five functions are the geometric functions of the general rotational surface of hyperbolic type $\mathcal{M}_2$ in the sense of [2]. The other three geometric functions of $\mathcal{M}_2$ are equal to zero.

Similarly to the elliptic case we obtain the following expressions for the Gauss curvature K, the curvature of the normal connection $\varkappa$, and the mean curvature vector field H of $\mathcal{M}_2$:

$$K = -\frac{\alpha^2\beta^2(fg'-f'g)^2(f'^2+g'^2) + (\alpha^2 fg' - \beta^2 f'g)(f''g'-f'g'')(\alpha^2 f^2+\beta^2 g^2)}{(f'^2+g'^2)^2(\alpha^2 f^2+\beta^2 g^2)^2};$$

$$\varkappa = \frac{\alpha\beta(fg'-f'g)\left((\alpha^2 f^2+\beta^2 g^2)(f''g'-f'g'') + (f'^2+g'^2)(\alpha^2 fg' - \beta^2 gf')\right)}{(f'^2+g'^2)^2(\alpha^2 f^2+\beta^2 g^2)^2};$$

$$H = \frac{(f'^2+g'^2)(\beta^2 f'g - \alpha^2 fg') + (\alpha^2 f^2+\beta^2 g^2)(f''g'-f'g'')}{2\,(f'^2+g'^2)^{\frac{3}{2}}\,(\alpha^2 f^2+\beta^2 g^2)}\,n_1.$$

Note that in the case $H \neq 0$ the normalized mean curvature vector field of a general rotational surface of hyperbolic type with plane meridian curves is timelike. So, we can formulate the following statement:

Proposition 3.4. *There are no quasi-minimal general rotational surfaces of hyperbolic type with plane meridian curves.*

If a Lorentz surface in $\mathbb{E}_2^4$ is neither minimal nor quasi-minimal, then it is worth considering the allied mean curvature vector field of the surface. The notion of allied mean curvature vector field of a Lorentz surface M_1^2 in $\mathbb{E}_2^4$ is defined by the formula

$$a(H) = \frac{\|H\|}{2} \operatorname{tr}(A_1 \circ A_2)\, n_2,$$

where $\{n_1 = \dfrac{H}{\|H\|}, n_2\}$ is an orthonormal base of the normal space of M_1^2; A_1 and A_2 are the shape operators corresponding to n_1 and n_2, respectively. M_1^2 is said to be a *Chen surface* (*Chen submanifold* in $\mathbb{E}_2^4$) if $a(H)$ vanishes identically [5].

Using formulas (10) and (12) for the general rotational surface $\mathcal{M}_1$ and $\mathcal{M}_2$, respectively, we can easily calculate that $a(H) = 0$. So, we get the following

Proposition 3.5. *Each general rotational surface of elliptic or hyperbolic type in $\mathbb{E}_2^4$ is a Chen surface.*

General rotational surfaces with plane meridian curves and pointwise 1-type Gauss map in $\mathbb{E}_2^4$ are studied in [1] under the assumption $\alpha = \beta = 1$ and $K = 0$. In [3], the classification of general rotational surfaces having zero mean curvature and pointwise 1-type Gauss map of second kind is given.

A special class of rotational surfaces with constant mean curvature in $\mathbb{E}_2^4$ is studied in [12]. These are general rotational surfaces with plane meridian curves in the case $\alpha = \beta = 1$ and $f(u) = \varphi(u)\sinh u$; $g(u) = \varphi(u)\cosh u$ for some smooth function $\varphi(u)$.

In what follows we study some basic classes of general rotational surfaces of elliptic and hyperbolic type in the case of arbitrary constants $\alpha > 0$ and $\beta > 0$. We give the classification of minimal general rotational surfaces, general rotational surfaces with parallel normalized mean curvature vector field, and flat general rotational surfaces. We describe analytically the class of general rotational surfaces with flat normal connection.

4. Minimal general rotational surfaces of elliptic or hyperbolic type

The study of minimal surfaces is one of the main topics in classical differential geometry. Recall that a surface in $\mathbb{E}_2^4$ is minimal if the mean curvature vector field $H = 0$. Each plane is a trivial minimal surface. We study only surfaces which do not contain any open part of a plane.

In the next theorem we classify all minimal general rotational surfaces of elliptic type.

Theorem 4.1. *Let $\mathcal{M}_1$ be a general rotational surface of elliptic type, defined by* (3)*. Then $\mathcal{M}_1$ is minimal if and only if the meridian curve m is determined by one of the following:*

(i) $f = cg^{\pm\frac{\alpha}{\beta}}$, $c = const$, $c \neq 0$, $\alpha \neq \beta$;

(ii) $\arcsin\left(\dfrac{\alpha f}{\sqrt{A}}\right) = \pm\dfrac{\alpha}{\beta}\arcsin\left(\dfrac{\beta g}{\sqrt{A}}\right) + C$, *where* $C = const$, $A = const$, $A > 0$, $\alpha \neq \beta$;

(iii) $(f+g)^2 = a\,(f-g)^2 + b$, *where* $a = const$, $a \neq 0$, $b = const$, *and* $\alpha = \beta$.

Proof: The mean curvature vector field H of a general rotational surface of elliptic type is given by (8). Hence, the surface $\mathcal{M}_1$ is minimal if and only if

$$\frac{g'f'' - f'g''}{f'^2 - g'^2} = \frac{\beta^2 g f' - \alpha^2 f g'}{\beta^2 g^2 - \alpha^2 f^2}.$$

On the other hand, from (10) it follows that the mean curvature vector field H is expressed as $H = \dfrac{\nu_2 - \nu_1}{2}\, n_2$, where ν_1 and ν_2 are given in (9). So, the condition $H = 0$ is equivalent to $\nu_1 = \nu_2$.

Now, let $\mathcal{M}_1$ is minimal, i.e. the equality $\nu_1 = \nu_2$ holds. Taking into account that $R' = 0$ and using derivative formulas (10) we obtain the equalities

$$\begin{aligned} 2\mu\gamma_2 - \nu_1\beta_2 &= x(\mu);\\ \mu\beta_2 - 2\nu_1\gamma_2 &= -x(\nu_1). \end{aligned} \tag{14}$$

If we assume that $\mu = 0$, from (9) we get $fg' - gf' = 0$, which implies $f = cg$, $c = const$. Straightforward computations show that in this case $\nu_1 = \nu_2 = 0$ and hence $\sigma(x, x) = \sigma(x, y) = \sigma(y, y) = 0$. So, the surface is totally geodesic, i.e. $\mathcal{M}_1$ is locally a plane.

So, further we consider $\mu \neq 0$. If we assume that $\nu_1 = \nu_2 = 0$, from (9) we get $g'f'' - f'g'' = 0$, which implies

$$\frac{f''}{f'} = \frac{g''}{g'}.$$

The solution of this equation is given by $f = ag + b$, $a = const \neq 0$, $b = const$. Without loss of generality we can consider $f(u)$ and $g(u)$ as follows:

$$f(u) = au + b; \quad g(u) = u. \tag{15}$$

In this case the function μ is expressed as $\mu = \dfrac{\alpha\beta\, b}{\sqrt{(a^2-1)}(\beta^2u^2 - \alpha^2(au+b)^2)}$. On the other hand, using (14) in the case $\nu_1 = 0$, we obtain $\mu\,\beta_2 = 0$. Since $\mu \neq 0$, we get $\beta_2 = 0$. The expression of β_2 in (9) implies $ff' - gg' = 0$. The last equality together with (15) gives $a^2 - 1 = 0$, which contradicts the assumption $f'^2 - g'^2 > 0$.

So, further we consider $\mu \neq 0$ and $\nu_1 = \nu_2 \neq 0$.

In the case $\alpha = \beta$, taking into account that $\nu_1 = \nu_2$, from (9) we get the equation:

$$\frac{f''g' - f'g''}{f'^2 - g'^2} = \frac{fg' - f'g}{f^2 - g^2}. \tag{16}$$

We denote $\phi(u) = \dfrac{f(u)}{g(u)}$ and $\psi(u) = \dfrac{f'(u)}{g'(u)}$. Then equation (16) is written in the form

$$\frac{\phi'(u)}{1 - \phi^2(u)} = \frac{\psi'(u)}{\psi^2(u) - 1},$$

which implies the following equality

$$\ln\left|\frac{1+\phi}{1-\phi}\right| = \ln\left|\frac{\psi - 1}{\psi + 1}\right| + const.$$

Hence, we obtain

$$\frac{(1+\phi)(1+\psi)}{(1-\phi)(1-\psi)} = const.$$

Now, using the expressions of $\phi(u)$ and $\psi(u)$ we get

$$\left((f+g)^2\right)' = a\left((f-g)^2\right)', \quad a = const \neq 0.$$

Hence, in the case $\alpha = \beta$ the relation between the functions f and g is:

$$(f+g)^2 = a\,(f-g)^2 + b,$$

where $a = const \neq 0$, $b = const$. Thus we obtain case (iii) in the statement of the theorem.

Further we consider the case $\alpha \neq \beta$. If $\mu^2 - \nu^2 = 0$, or equivalently $\nu_1 = \nu_2 = \pm\mu$, then the expressions of ν_2 and μ in (9) give the equation:

$$\beta^2 g f' - \alpha^2 f g' = \pm\alpha\beta(fg' - gf'),$$

which can be written in the form:

$$\frac{f'}{f} = \pm\frac{\alpha}{\beta}\frac{g'}{g}.$$

The solution of the last equation is:

$$f = cg^{\pm\frac{\alpha}{\beta}}, \quad c = const,\ c \neq 0,$$

which gives (i) in the statement of the theorem.

Now we consider the case $\mu^2 - \nu^2 \neq 0$. Denote $\nu := \nu_1 = \nu_2$. Using equalities (14) we get the equations:

$$x(\mu) = 2\mu\gamma_2 - \nu\beta_2;$$
$$x(\nu) = 2\nu\gamma_2 - \mu\beta_2,$$

which imply

$$x(\mu^2 - \nu^2) = 4(\mu^2 - \nu^2)\gamma_2. \tag{17}$$

Formula (17) together with the equality $\gamma_2 = -x(\ln(\sqrt{-G})$ give us

$$x(\ln|\mu^2 - \nu^2|) = -4x(\ln(\sqrt{-G}),$$

which implies

$$x(\ln|(\mu^2 - \nu^2)G^2|) = 0.$$

Since the functions μ, ν, and G do not depend on the parameter v, we obtain

$$|(\mu^2 - \nu^2)G^2| = c^2, \quad c = const \neq 0.$$

Using the expressions of μ and ν in (9) and taking into account $f'^2 - g'^2 = 1$, we obtain the equation:

$$\alpha^2\beta^2(fg' - f'g)^2 - (\beta^2 f'g - \alpha^2 fg')^2 = c^2,$$

or equivalently:

$$(\alpha^2 - \beta^2)(\beta^2 f'^2 g^2 - \alpha^2 f^2(f'^2 - 1)) = c^2.$$

Since $\alpha^2 - \beta^2 \neq 0$, putting $A = \dfrac{c^2}{\alpha^2 - \beta^2}$, we obtain the following expression for f'^2:

$$f'^2 = \frac{A - \alpha^2 f^2}{\beta^2 g^2 - \alpha^2 f^2}. \tag{18}$$

Using that $g'^2 = f'^2 - 1$ we get:

$$g'^2 = \frac{A - \beta^2 g^2}{\beta^2 g^2 - \alpha^2 f^2}. \tag{19}$$

Since $f'^2 > 0$ and $g'^2 > 0$, we get $A > \alpha^2 f^2$ and $A > \beta^2 g^2$, so $A > 0$, i.e. $\alpha^2 > \beta^2$. Equations (18) and (19) imply

$$\frac{f'^2}{A - \alpha^2 f^2} = \frac{g'^2}{A - \beta^2 g^2}.$$

The last equality gives the following relation between the functions f and g:

$$\arcsin\left(\frac{\alpha f}{\sqrt{A}}\right) = \pm\frac{\alpha}{\beta}\arcsin\left(\frac{\beta g}{\sqrt{A}}\right) + C, \quad C = const.$$

This corresponds to case (ii) in the statement of the theorem.

Conversely, if one of (i), (ii), or (iii) holds, then by straightforward computations it follows that $\nu_1 = \nu_2$, i.e. $\mathcal{M}_1$ is a minimal surface. □

The following theorem gives the classification of all minimal general rotational surfaces of hyperbolic type.

Theorem 4.2. *Let $\mathcal{M}_2$ be a general rotational surface of hyperbolic type, defined by* (11). *Then $\mathcal{M}_2$ is minimal if and only if the meridian curve m is determined by one of the following:*

(i) $f = cg^{\mp\frac{\alpha}{\beta}}, \quad c = const,\ c \neq 0,\ \alpha \neq \beta;$

(ii) $\alpha f + \sqrt{\alpha^2 f^2 - A} = C\left(\beta g + \sqrt{\beta^2 g^2 + A}\right)^{\pm\frac{\alpha}{\beta}},\ C = const, A = const, AC \neq 0, \alpha \neq \beta;$

(iii) $\arctan\left(\dfrac{f'}{g'}\right) = -\arctan\left(\dfrac{f}{g}\right) + c, \quad c = const,\ and\ \alpha = \beta.$

Proof: Similarly to the proof of Theorem 4.1, we obtain that $\mu \neq 0$ and $\nu_1 = \nu_2 \neq 0$. We denote $\nu := \nu_1 = \nu_2$.

In the case $\alpha = \beta$, using that $\nu_1 = \nu_2$, from (13) we get the equation

$$\frac{f''g' - f'g''}{f'^2 + g'^2} = \frac{fg' - f'g}{f^2 + g^2},$$

which implies

$$\left(\arctan\left(\frac{f'}{g'}\right)\right)' = -\left(\arctan\left(\frac{f}{g}\right)\right)'.$$

So, the relation between the functions f and g is given by:

$$\arctan\left(\frac{f'}{g'}\right) = -\arctan\left(\frac{f}{g}\right) + c, \quad c = const,$$

which corresponds to case (iii) of the theorem.

Now, we consider the case $\alpha \neq \beta$. If $\mu^2 - \nu^2 = 0$, then as in the elliptic case we get the equation:

$$\alpha^2 fg' - \beta^2 gf' = \pm\alpha\beta(fg' - gf'),$$

whose solution is:

$$f = cg^{\mp\frac{\alpha}{\beta}}, \quad c = const, c \neq 0.$$

This is case (i) in the statement of the theorem.

If $\mu^2 - \nu^2 \neq 0$, then from $R' = 0$ and (12) we obtain the equation

$$x(\mu^2 - \nu^2) = 4(\mu^2 - \nu^2)\gamma_2.$$

On the other hand, we have $\gamma_2 = -x(\ln(\sqrt{-G})$, so we get

$$|\mu^2 - \nu^2|G^2 = c^2, \quad c = const \neq 0.$$

Without loss of generality we assume that $f'^2 + g'^2 = 1$. So, we obtain the equation

$$(\beta^2 f'^2 g^2 - \alpha^2 f^2(1 - f'^2))(\alpha^2 - \beta^2) = c^2.$$

Since $\alpha^2 - \beta^2 \neq 0$, putting $A = \dfrac{c^2}{\alpha^2 - \beta^2}$, we get the following expressions for f'^2 and g'^2:

$$f'^2 = \frac{\alpha^2 f^2 - A}{\alpha^2 f^2 + \beta^2 g^2}, \quad g'^2 = \frac{\beta^2 g^2 + A}{\alpha^2 f^2 + \beta^2 g^2},$$

which imply

$$\frac{f'^2}{\alpha^2 f^2 - A} = \frac{g'^2}{\beta^2 g^2 + A}. \tag{20}$$

Note that $\alpha^2 f^2 - A > 0$ and $\beta^2 g^2 + A > 0$, since $f'^2 > 0$ and $g'^2 > 0$. Equation (20) implies that the functions f and g, which describe the meridian curve m of $\mathcal{M}_2$, satisfy the following relation:

$$\ln\left|\alpha f + \sqrt{\alpha^2 f^2 - A}\right| = \pm\frac{\alpha}{\beta}\ln\left|\beta g + \sqrt{\beta^2 g^2 + A}\right| + c,$$

where $c = const$. Finally, we obtain

$$\alpha f + \sqrt{\alpha^2 f^2 - A} = C\left(\beta g + \sqrt{\beta^2 g^2 + A}\right)^{\pm\frac{\alpha}{\beta}}.$$

This is case (ii) in the theorem.

□

5. General rotational surfaces of elliptic or hyperbolic type with parallel normalized mean curvature vector field

In the case $H \neq 0$ the normalized mean curvature vector field of the rotational surface of elliptic type $\mathcal{M}_1$ is n_2. So, $\mathcal{M}_1$ has parallel normalized mean curvature vector field if and only if $D_x n_2 = D_y n_2 = 0$. The next theorem describes all general rotational surfaces of elliptic type with parallel normalized mean curvature vector field.

Theorem 5.1. *Let $\mathcal{M}_1$ be a general rotational surface of elliptic type, defined by* (3)*. Then $\mathcal{M}_1$ has parallel normalized mean curvature vector field if and only if the meridian curve m is determined by*

$$f(u) = \pm\sqrt{u^2 - C^2}; \quad g(u) = u, \quad C = const \neq 0.$$

Proof: It follows from (5) that $D_x n_2 = D_y n_2 = 0$ if and only if the functions f and g satisfy the following differential equation:

$$ff' - gg' = 0,$$

which implies that $f^2 = g^2 + C_1$ for some constant C_1. Without loss of generality we can assume that $g(u) = u$. Then $f(u) = \pm\sqrt{u^2 + C_1}$. Since $f'^2 - g'^2 > 0$, we obtain $C_1 < 0$. Hence $f(u) = \pm\sqrt{u^2 - C^2}$ for some constant $C \neq 0$.

□

Using the last theorem and equality (8) we obtain that the mean curvature vector field H of a general rotational surface of elliptic type with parallel normalized mean curvature vector field is given by:

$$H = \frac{1}{C}\, n_2.$$

Hence, the condition on $\mathcal{M}_1$ to have parallel normalized mean curvature vector field implies $\langle H, H\rangle = -\dfrac{1}{C^2} = const$. So, the following statement holds:

Corollary 5.2. *If $\mathcal{M}_1$ has parallel normalized mean curvature vector field, then it has parallel mean curvature vector field.*

The classification of general rotational surfaces of hyperbolic type with parallel normalized mean curvature vector field is given in the next theorem.

Theorem 5.3. *Let $\mathcal{M}_2$ be a general rotational surface of hyperbolic type, defined by* (11)*. Then $\mathcal{M}_2$ has parallel normalized mean curvature vector field if and only if the meridian curve m is determined by*

$$f(u) = \pm\sqrt{C^2 - u^2}; \quad g(u) = u, \quad C = const \neq 0.$$

Proof: In the case $H \neq 0$ the normalized mean curvature vector field of $\mathcal{M}_2$ is n_1. Using (12) and (13), we get that $\mathcal{M}_2$ has parallel normalized mean curvature vector field if and only if the functions f and g satisfy the equation

$$ff' + gg' = 0,$$

which implies $f^2 + g^2 = C^2$ for some constant $C \neq 0$. Without loss of generality we can assume that $g(u) = u$ and then the function $f(u)$ has the form $f(u) = \pm\sqrt{C^2 - u^2}$.

□

Calculating the mean curvature vector field H of a general rotational surface of hyperbolic type with parallel normalized mean curvature vector field, we obtain

$$H = \mp\frac{1}{C}\, n_1.$$

The last equality shows that $\|H\| = const$, so the following statement holds true:

Corollary 5.4. *If $\mathcal{M}_2$ has parallel normalized mean curvature vector field, then it has parallel mean curvature vector field.*

6. Flat general rotational surfaces of elliptic or hyperbolic type

In this section we give the classification of flat general rotational surfaces of elliptic or hyperbolic type.

Theorem 6.1. *Let $\mathcal{M}_1$ be a general rotational surface of elliptic type, defined by* (3)*. Then $\mathcal{M}_1$ is flat if and only if the meridian curve m is determined by one of the following:*

(i) $\beta^2 g^2 - \alpha^2 f^2 = a^2(u + c)^2$, where $a = const \neq 0$, $c = const$;

(ii) $\alpha^2 f^2 - \beta^2 g^2 = C$, where $C = const$, $C < 0$.

Proof: A Lorentz surface is flat, if its Gauss curvature is zero. In terms of the functions μ, ν_1 and ν_2 participating in formulas (10), the condition on $\mathcal{M}_1$ to have zero Gauss curvature is expressed as $\mu^2 + \nu_1\nu_2 = 0$.

Now, let $\mu^2 + \nu_1\nu_2 = 0$, i.e. $K = 0$. Using that $R' = 0$ from derivative formulas (10) we obtain

$$x(\gamma_2) = (\gamma_2)^2. \tag{21}$$

Without loss of generality we can assume that $f'^2 - g'^2 = 1$. Then $x = \dfrac{\partial}{\partial u}$.

If $\gamma_2 \neq 0$ from equality (21) we get the equation :

$$\frac{\gamma_2'}{(\gamma_2)^2} = 1,$$

whose solution is

$$\gamma_2(u) = -\frac{1}{u + c}, \quad c = const.$$

On the other hand, $\gamma_2 = -x(\ln\sqrt{\beta^2 g^2 - \alpha^2 f^2})$. Consequently,

$$(\ln\sqrt{\beta^2 g^2 - \alpha^2 f^2})' = \frac{1}{u+c}.$$

The last equation gives the following relation between the functions f and g determining the meridian curve m:

$$\beta^2 g^2 - \alpha^2 f^2 = a^2(u+c)^2,$$

where $a = const \neq 0$ and $c = const$.

If $\gamma_2 = 0$, from (9) we get the equation

$$\alpha^2 f f' - \beta^2 g g' = 0,$$

which implies

$$\alpha^2 f^2 - \beta^2 g^2 = C, \quad C = const.$$

Since we consider surfaces with $\alpha^2 f^2 - \beta^2 g^2 < 0$, so the constant C is negative.

Conversely, if (i) or (ii) holds, then by straightforward computations we get $\mu^2 + \nu_1\nu_2 = 0$, i.e. the surface $\mathcal{M}_1$ is flat.

□

The next theorem describes flat general rotational surfaces of hyperbolic type.

Theorem 6.2. *Let $\mathcal{M}_2$ be a general rotational surface of hyperbolic type, defined by* (11). *Then $\mathcal{M}_2$ is flat if and only if the meridian curve m is determined by one of the following:*

(i) $\alpha^2 f^2 + \beta^2 g^2 = a^2(u+c)^2, \quad a = const \neq 0,\ c = const;$

(ii) $\alpha^2 f^2 + \beta^2 g^2 = C, \quad C = const.$

Proof: The proof in the hyperbolic case is similar to the elliptic one. Again we have that a general rotational surface of hyperbolic type is flat if and only if the functions μ, ν_1 and ν_2 participating in (12) satisfy the condition $\mu^2 + \nu_1\nu_2 = 0$. The function γ_2 satisfies the equation

$$x(\gamma_2) = (\gamma_2)^2.$$

In the case $\gamma_2 \neq 0$, assuming that $f'^2 + g'^2 = 1$, we obtain:

$$\gamma_2(u) = -\frac{1}{u+c}, \quad c = const.$$

On the other hand, $\gamma_2 = -x(\ln\sqrt{\alpha^2 f^2 + \beta^2 g^2})$, so

$$(\ln\sqrt{\alpha^2 f^2 + \beta^2 g^2})' = \frac{1}{u+c}.$$

The solution of the last differential equation is:

$$\alpha^2 f^2 + \beta^2 g^2 = a^2(u+c)^2,$$

where $a = const \neq 0$, $c = const$.

If $\gamma_2 = 0$ we obtain

$$\alpha^2 f f' + \beta^2 g g' = 0,$$

or equivalently

$$\alpha^2 f^2 + \beta^2 g^2 = C, \quad C = const.$$

□

7. General rotational surfaces of elliptic or hyperbolic type with flat normal connection

A surface is said to have flat normal connection if the curvature of the normal connection is zero. Each surface with parallel normalized mean curvature vector field has flat normal connection. In Section 5, we classified all general rotational surfaces with parallel normalized mean curvature vector field. Note that according to Corollary 5.2 and Corollary 5.4, in the class of general rotational surfaces of both elliptic and hyperbolic type the condition of parallel normalized mean curvature vector field is equivalent to the condition of parallel mean curvature vector field. So, here we shall consider general rotational surfaces with flat normal connection but non-parallel mean curvature vector field, i.e. we assume that $\beta_2 \neq 0$.

Theorem 7.1. *Let $\mathcal{M}_1$ be a general rotational surface of elliptic type, defined by* (3)*. Then $\mathcal{M}_1$ has flat normal connection and non-parallel mean curvature vector field if and only if the meridian curve m is determined by one of the following:*

(i) $f = c\,g$, where $c = const$, $1 < c^2 < \dfrac{\beta^2}{\alpha^2}$, and $\alpha < \beta$. In this case, $\mathcal{M}_1$ is a developable ruled surface in $\mathbb{E}^4_2$;

(ii) $$\frac{ff' - gg'}{\sqrt{f'^2 - g'^2}\sqrt{\beta^2 g^2 - \alpha^2 f^2}} = C, \text{ where } C = const,\ C \neq 0.$$

Proof: The curvature of the normal connection of the general rotational surface $\mathcal{M}_1$ is given by formula (7). Hence, $\mathcal{M}_1$ has flat normal connection if and only if

$$(fg' - gf')\left((\beta^2 g^2 - \alpha^2 f^2)(g'f'' - f'g'') + (f'^2 - g'^2)(\beta^2 gf' - \alpha^2 fg')\right) = 0.$$

The last equation leads to the following two cases:

(i) $fg' - gf' = 0$, i.e. $f = c\,g$ for some constant $c \neq 0$. Since we consider surfaces for which $f'^2 - g'^2 > 0$ and $\alpha^2 f^2 - \beta^2 g^2 < 0$, so the constant c satisfies $1 < c^2 < \dfrac{\beta^2}{\alpha^2}$, and hence $\alpha < \beta$. By straightforward computations we get $\nu_1 = 0$, $\nu_2 = \dfrac{c(\beta^2 - \alpha^2)}{g\sqrt{c^2 - 1}(\beta^2 - c^2\alpha^2)} \neq 0$, $\mu = 0$, $\gamma_2 = -\dfrac{1}{g\sqrt{c^2 - 1}} \neq 0$, $\beta_2 = \dfrac{\alpha\beta\sqrt{c^2 - 1}}{g(\beta^2 - c^2\alpha^2)} \neq 0$. In this case $\mathcal{M}_1$ is a ruled surface, since the meridian curve m is a straight line. Using formulas (10) we obtain that $\nabla'_x n_1 = 0$, $\nabla'_x n_2 = 0$, so, the normal space is constant at each point of a fixed straight line. Consequently, $\mathcal{M}_1$ is a developable ruled surface in $\mathbb{E}^4_2$.

(ii) $fg' - gf' \neq 0$. In this case the functions f and g satisfy the equation

$$(\beta^2 g^2 - \alpha^2 f^2)(g'f'' - f'g'') + (f'^2 - g'^2)(\beta^2 gf' - \alpha^2 fg') = 0,$$

which is equivalent to $\nu_1 + \nu_2 = 0$. Using that $R' = 0$, from derivative formulas (10) under the assumption $\nu_1 + \nu_2 = 0$, we obtain $\gamma_2\beta_2 = x(\beta_2)$. Since $\beta_2 \neq 0$, we get the equation

$$\frac{x(\beta_2)}{\beta_2} = \gamma_2,$$

which implies $x(\ln|\beta_2|) = \gamma_2$. On the other hand, $\gamma_2 = -x(\ln\sqrt{-G})$. Consequently,

$$x\left(\ln|\beta_2\sqrt{-G}|\right) = 0. \tag{22}$$

Now, using that the functions β_2 and G do not depend on the parameter v, from (22) it follows that

$$\beta_2\sqrt{-G} = const.$$

Having in mind the expression of β_2 given in (9), we obtain that the functions f and g satisfy the following differential equation

$$\frac{ff' - gg'}{\sqrt{f'^2 - g'^2}\sqrt{\beta^2 g^2 - \alpha^2 f^2}} = C,$$

where $C = const$, $C \neq 0$.

□

Similarly to the proof of Theorem 7.1 we obtain the following characterization of general rotational surfaces of hyperbolic type with flat normal connection.

Theorem 7.2. *Let $\mathcal{M}_2$ be a general rotational surface of hyperbolic type, defined by* (11). *Then $\mathcal{M}_2$ has flat normal connection and non-parallel mean curvature vector field if and only if the meridian curve m is determined by one of the following:*

(i) $f = c\,g$, where $c = const$, $c \neq 0$, and $\alpha \neq \beta$. In this case, $\mathcal{M}_1$ is a developable ruled surface in $\mathbb{E}_2^4$;

(ii) $\dfrac{ff' + gg'}{\sqrt{f'^2 + g'^2}\sqrt{\alpha^2 f^2 + \beta^2 g^2}} = C$, *where $C = const$, $C \neq 0$.*

Acknowledgments: The second author is partially supported by the Bulgarian National Science Fund, Ministry of Education and Science of Bulgaria under contract DFNI-I 02/14. The third author is supported by TÜBİTAK (Project Name: Y_EUCL2TIP, Project Number: 114F199).

This work was done during the third author's visit at the Institute of Mathematics and Informatics, Bulgarian Academy of Sciences in June 2015.

References

[1] Aksoyak F., Yayli Y., *General rotational surfaces with pointwise 1-type Gauss map in pseudo-Euclidean space $\mathbb{E}_2^4$*. Indian J. Pure Appl. Math., **46**, no. 1 (2015), 107–118.

[2] Aleksieva Y., Ganchev G., Milousheva V., *On the theory of Lorentz surfaces with parallel normalized mean curvature vector field in pseudo-Euclidean 4-space*. J. Korean Math. Soc., **53**, (2016).

[3] Bektaş B., Canfes E., Dursun U., *On rotational surfaces in pseudo-Euclidean space $\mathbb{E}_t^4$ with pointwise 1-type Gauss map*. ArXiv 1508.03294v1.

[4] Chen B.-Y., *Surfaces with parallel normalized mean curvature vector*. Monatsh. Math., **90**, no. 3 (1980), 185–194.

[5] Chen B.-Y., *Pseudo-Riemannian geometry, δ-invariants and applications*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2011.

[6] Dursun U., *On spacelike rotational surfaces with pointwise 1-type Gauss map*. Bull. Korean Math. Soc., **52**, no. 1 (2015), 301–312.

[7] Dursun U., Turgay N., *General rotational surfaces in Euclidean space $\mathbb{E}^4$ with pointwise 1-type Gauss map*. Math. Commun., **17** (2012), 71–81.

[8] Dursun U., Turgay N., *Minimal and pseudo-umbilical rotational surfaces in Euclidean space $\mathbb{E}^4$*. Mediterr. J. Math. **10**, no. 1 (2013), 497–506.

[9] Ganchev G., Milousheva V., *Invariants of lines on surfaces in $\mathbb{R}^4$*. C. R. Acad. Bulgare Sci., **63**, no. 6 (2010), 835-842.

[10] Ganchev G., Milousheva V., *Quasi-minimal rotational surfaces in pseudo-Euclidean four-dimensional space*. Cent. Eur. J. Math., **12**, no. 10 (2014), 1586–1601.

[11] Ganchev G., Milousheva V., *General rotational surfaces in the 4-dimensional Minkowski space*. Turk. J. Math., **38**, no. 5 (2014), 883–895.

[12] HuiLi L., GuiLi L., *Rotation surfaces with constant mean curvature in 4-dimensional pseudo-Euclidean space*. Kyushu J. Math., **48**, no. 1 (1994), 35–42.

[13] Milousheva V., *General rotational surfaces in* $\mathbb{R}^4$ *with meridians lying in two-dimensional planes*. C. R. Acad. Bulgare Sci., **63**, no. 3 (2010), 339–348.
[14] Moore C., *Surfaces of rotation in a space of four dimensions*. Ann. of Math., 2nd Ser., **21**, no. 2 (1919), 81-93.
[15] Moore C., *Rotation surfaces of constant curvature in space of four dimensions*. Bull. Amer. Math. Soc., **26**, no. 10 (1920), 454-460.
[16] Rosca R., *On null hypersurfaces of a Lorentzian manifold*. Tensor (N.S.) **23** (1972), 66–74.

Faculty of Mathematics and Informatics, Sofia University, 5 James Bourchier blvd., 1164 Sofia, Bulgaria
E-mail address: yana_a_n@fmi.uni-sofia.bg

Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, Acad. G. Bonchev Str. bl. 8, 1113, Sofia, Bulgaria; "L. Karavelov" Civil Engineering Higher School, 175 Suhodolska Str., 1373 Sofia, Bulgaria
E-mail address: vmil@math.bas.bg

Istanbul Technical University, Faculty of Science and Letters, Department of Mathematics, 34469 Maslak, Istanbul, Turkey
E-mail address: turgayn@itu.edu.tr